

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Pearson Edexcel
Level 3 GCE

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

Thursday 20 June 2019

Morning (Time: 1 hour 30 minutes)

Paper Reference **9FM0/3C**

Further Mathematics

Advanced

Paper 3C: Further Mechanics 1

You must have:

Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

--

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for algebraic manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Unless otherwise indicated, whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$ and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P62674A

©2019 Pearson Education Ltd.

1/1/1



Pearson

3. A particle P , of mass 0.5 kg , is moving with velocity $(4\mathbf{i} + 4\mathbf{j}) \text{ m s}^{-1}$ when it receives an impulse \mathbf{I} of magnitude 2.5 N s .

As a result of the impulse, the direction of motion of P is deflected through an angle of 45°

Given that $\mathbf{I} = (\lambda\mathbf{i} + \mu\mathbf{j}) \text{ N s}$, find all the possible pairs of values of λ and μ .

(9)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



4. A car of mass 600 kg pulls a trailer of mass 150 kg along a straight horizontal road. The trailer is connected to the car by a light inextensible towbar, which is parallel to the direction of motion of the car. The resistance to the motion of the trailer is modelled as a constant force of magnitude 200 N. At the instant when the speed of the car is $v \text{ ms}^{-1}$, the resistance to the motion of the car is modelled as a force of magnitude $(200 + \lambda v) \text{ N}$, where λ is a constant.

When the engine of the car is working at a constant rate of 15 kW, the car is moving at a constant speed of 25 ms^{-1}

- (a) Show that $\lambda = 8$

(4)

Later on, the car is pulling the trailer up a straight road inclined at an angle θ to the horizontal, where $\sin \theta = \frac{1}{15}$

The resistance to the motion of the trailer from non-gravitational forces is modelled as a constant force of magnitude 200 N at all times. At the instant when the speed of the car is $v \text{ ms}^{-1}$, the resistance to the motion of the car from non-gravitational forces is modelled as a force of magnitude $(200 + 8v) \text{ N}$.

The engine of the car is again working at a constant rate of 15 kW.

When $v = 10$, the towbar breaks. The trailer comes to instantaneous rest after moving a distance d metres up the road from the point where the towbar broke.

- (b) Find the acceleration of the car immediately after the towbar breaks.

(4)

- (c) Use the work-energy principle to find the value of d .

(4)



5. A particle P of mass $3m$ and a particle Q of mass $2m$ are moving along the same straight line on a smooth horizontal plane. The particles are moving in opposite directions towards each other and collide directly.

Immediately before the collision the speed of P is u and the speed of Q is $2u$.

Immediately after the collision P and Q are moving in opposite directions.

The coefficient of restitution between P and Q is e .

- (a) Find the range of possible values of e , justifying your answer.

(8)

Given that Q loses 75% of its kinetic energy as a result of the collision,

- (b) find the value of e .

(3)



6. [In this question \mathbf{i} and \mathbf{j} are perpendicular unit vectors in a horizontal plane.]

A smooth uniform sphere A has mass 0.2 kg and another smooth uniform sphere B , with the same radius as A , has mass 0.4 kg .

The spheres are moving on a smooth horizontal surface when they collide obliquely. Immediately before the collision, the velocity of A is $(3\mathbf{i} + 2\mathbf{j})\text{ m s}^{-1}$ and the velocity of B is $(-4\mathbf{i} - \mathbf{j})\text{ m s}^{-1}$

At the instant of collision, the line joining the centres of the spheres is parallel to \mathbf{i}

The coefficient of restitution between the spheres is $\frac{3}{7}$

- (a) Find the velocity of A immediately after the collision. (7)
- (b) Find the magnitude of the impulse received by A in the collision. (2)
- (c) Find, to the nearest degree, the size of the angle through which the direction of motion of A is deflected as a result of the collision. (3)



7. A particle P , of mass m , is attached to one end of a light elastic spring of natural length a and modulus of elasticity kmg .

The other end of the spring is attached to a fixed point O on a ceiling.

The point A is vertically below O such that $OA = 3a$

The point B is vertically below O such that $OB = \frac{1}{2}a$

The particle is held at rest at A , then released and first comes to instantaneous rest at the point B .

(a) Show that $k = \frac{4}{3}$ (3)

(b) Find, in terms of g , the acceleration of P immediately after it is released from rest at A . (3)

(c) Find, in terms of g and a , the maximum speed attained by P as it moves from A to B . (6)



